Infant SAFE SLEP

- Always place your baby on his or her back to sleep, for naps and at night.
- Use a **firm sleep surface**, such as a mattress in a safety-approved crib, covered by a fitted sheet.
- Avoid wedges, positioners or other products unless prescribed by your baby's doctor.
- Keep pillows, bottles, toys, crib bumpers, and loose bedding out of your baby's sleep area.
- Don't sleep with your baby in a bed, on a chair or couch put your baby in his or her own bed.
- Keeping your **baby's sleep area in the same room where you sleep reduces the risk of SIDS** and other sleep-related causes of infant death.
- Offer your baby a pacifier that is not attached to a string for naps and at night. If your baby is breastfed, wait until your baby is one month old before offering a pacifier.
- **Keep your baby warm, but not hot.** Dress your baby in one layer of clothing extra than what you would wear to be comfortable and leave the blanket out of the crib.
- Follow your health care provider's guidance on your baby's vaccines and regular health checkups. Talk with your doctor if you have any questions about how your baby sleeps.
- Give your baby plenty of time on his or her tummy when awake and when someone is watching.
- Do not smoke or allow smoking around your baby.
- Place your **baby's crib away from curtains or blinds** to avoid strangulation by cords.
- Make sure anyone caring for your baby knows about safe sleep practices.

*The actions listed here are based on recommendations from the American Academy of Pediatrics Task Force on SIDS.



Guidelipes



DSHS 22-1577 (10/14)