OICS know the facts

What are the risks of misusing prescription pain relievers (opioids)?

Talk with your health care provider about all the prescription drugs you take.

- Mental confusion
- Drowsiness
- Changes in brain that build tolerance, leading to dependence

WITHDRAWAL SYMPTOMS: Restlessness, muscle and bone pain, insomnia, diarrhea, vomiting, cold flashes with goose bumps, kicking movements, severe craving that can lead to relapse

TOTAL BODY EFFECTS: Euphoria, warm flushing, poor nutrition, fatal overdose

PREGNANCY AND INFANTS: Miscarriage, low birth weight, developmental delays, addiction in infants

..... O linfectious diseases such as HIV, Hepatitis B and Hepatitis C if injected

- Increased and/or decreased blood pressure
- ······ O Slow, irregular breathing that can be fatal
 - Constipation Cramping
 - Nausea
- ······ O HIV and other sexually transmitted infections from unsafe sex while intoxicated

Symptoms of overdose

COLD/CLAMMY SKIN DRY MOUTH WEAK PULSE DROWSINESS LOW OOD PRESSURE **COMA** BLUE NAILS AND LIPS MUSCLE SPASMS NON RESPONSIVE SEIZURES SMALL PUPILS NO BREATHING OR SHALLOW BREATHING

Need help?


For free referrals and emotional support: Washington Recovery

Help Line 1-866-789-1511

Teen Link

1-866-833-6546

Quick facts



- O Prescription opioids are medications that relieve
- They can be addictive if not used as directed by your doctor.
- More people die from opioid overdoses than from any other drug.



Nearly 17,000 people die each year from prescription pain reliever overdoses.

Lock up your meds and safely discard them. Visit: www.TakeBackYourMeds.org



